

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 7/50	A1	(11) International Publication Number: WO 94/05256 (43) International Publication Date: 17 March 1994 (17.03.94)
(21) International Application Number: PCT/US93/08137 (22) International Filing Date: 30 August 1993 (30.08.93) (30) Priority data: 07/941,594 8 September 1992 (08.09.92) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventor: KALLA, Karen, Kay ; 3453-1B Steeplechase Lane, Loveland, OH 45140 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).	(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: MILD, SUBSTANTIALLY COLORLESS SHAMPOO COMPOSITION		
(57) Abstract A mild, substantially colorless shampoo composition comprising: (a) from about 2 % to about 20 %, by weight, of an alkyl ethoxylated sulfate anionic surfactant having an average degree of ethoxylation of at least about 2.0; (b) from about 1 % to about 15 % of an amphoteric surfactant selected from the group consisting of betaine surfactant, imidazoline surfactants, aminoalkanoate surfactants, and iminodialkanoate surfactants, and mixtures thereof; (c) from about 0.5 % to about 10 %, by weight of an N-acyl amino acid surfactant; (d) at least about 0.05 %, by weight of the composition, of carbonyl-containing perfume compounds; and (e) water; wherein said shampoo composition is substantially free of alkyl sulfate anionic surfactant, free primary amines and ammonium ions, and amide foam boosters, and said shampoo composition has an absorbance value at 440 nm in a 1.0 cm path length of no greater than about 0.030 after storage at 38 °C for 30 days.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

-1-

MILD, SUBSTANTIALLY COLORLESS SHAMPOO COMPOSITION

TECHNICAL FIELD

5 The present invention is related to shampoo compositions. More specifically, the present invention relates to mild, substantially colorless shampoos, especially those that are useful for application to the hair.

BACKGROUND OF THE INVENTION

10 Mild shampoo compositions which provide low irritation to the skin are highly desirable. Conventional shampoos contain high levels of harsh anionic surfactants. These surfactants can provide excellent lathering and cleaning characteristics. However, they can also penetrate the skin and destroy its integrity.
15 This results, at the very least, in rough and/or dry skin, and can ultimately lead to red, irritated skin. Ideal shampoo compositions should provide sufficient lathering and cleaning benefits to cleanse the hair while at the same time causing little or no irritation to the skin. This is particularly essential for
20 shampoo compositions used on babies, small children, or adults with dry or sensitive skin.

Mild shampoo compositions, in general, are well known in the art having been taught in, for example, European Patent Application 0250181, published December 23, 1987; U.S. Patent 4,578,216,
25 Fujii et al., issued March 25, 1986; U.S. Patent 4,726,915, Verdicchio, issued February 23, 1988; U.K. Patent Specification 1,508,929, published April 26, 1978; European Patent Application 0160269, published November 6, 1985; U.S. Patent 4,435,300, Guth et al., issued March 6, 1984; U.S. Patent 4,426,310, Vernunica,
30 issued January 17, 1984; U.S. Patent 3,950,417, Verdicchio et al., issued April 13, 1976; U.S. Patent 4,443,362, Guth et al., issued April 17, 1984; U.S. Patent 4,654,207, Preston, issued March 31, 1987; U.S. Patent 4,851,154, Grollier et al., issued July 25, 1989; U.S. Patent 4,292,212, Melby, issued September 29, 1981; and
35 U.S. Patent 4,329,334, Su et al., issued May 11, 1982. These

- 2 -

references teach the use of a wide variety of surfactant combinations to achieve mildness. These surfactants include mild anionic surfactants, such as ethoxylated alkyl sulfates, amphoteric surfactants, nonionic surfactants, and various combinations thereof.

5 In addition to the physical effect harsh shampoos can have on the skin, consumers are also influenced by aesthetic factors relating to the shampoo. For example, high lathering is associated with good cleaning. Also, clear and colorless products are associated with purity and mildness. It would be highly
10 desirable to provide a mild shampoo which is both clear and colorless, and which can provide high levels of lather upon use.

Unfortunately, the formulation of mild shampoos is generally an exercise in trade-offs and compromises. Mild shampoos
15 conventionally have relatively poor lathering and are not colorless. Clear colorless shampoos tend to be harsh to the skin, to be characterized by poor lather and/or to suffer from discoloration after extended storage periods.

Hence, it is an object of the present invention to provide a
20 mild shampoo composition which is substantially colorless even over extended periods of storage, and which provides good lathering while at the same time providing good cleaning and being mild to the skin.

This and other objects will become readily apparent from the
25 detailed description which follows.

Unless otherwise indicated, all percentages are calculated by weight of the total composition and all ratios are calculated on a weight basis.

The compositions and processes herein can comprise, consist
30 essentially of, or consist of any of the critical and optional elements and steps described in this disclosure.

SUMMARY OF THE INVENTION

The present invention provides a mild, substantially colorless
shampoo composition that can provide good lathering
35 characteristics which comprises:

- 3 -

- (a) from about 2% to about 20%, by weight, of an alkyl ethoxylated sulfate anionic surfactant having an average degree of ethoxylation of at least about 2.0;
- 5 (b) from about 1% to about 15%, by weight, of an amphoteric surfactant selected from the group consisting of betaine surfactants, imidazoline surfactants, aminoalkanoate surfactants, and iminodialkanoate surfactants, and mixtures thereof;
- 10 (c) from about 0.5% to about 10%, by weight, of an N-acylamino acid surfactant, or salt thereof;
- (d) at least about 0.05%, by weight, of the composition, of carbonyl-containing perfume compounds; and
- (e) water;

15 wherein said shampoo composition is substantially free of alkyl sulfate anionic surfactant, primary amines, and ammonia, and said shampoo composition has an absorbance value at 440nm in a 1.0cm path length of no greater than about 0.030 after storage at 38°C for 30 days.

20 Although it is not intended to necessarily limit the invention by theory, it is believed that ammonia, which is typically introduced into the compositions in the form of ammonium ions, and/or primary amines react with carbonyl functionalities present in compounds found in most perfume formulations to form Schiff bases. These Schiff bases discolor the compositions. Insofar as

25 it is desired to use conventional perfume formulations in the present shampoo compositions, the compositions hereof contain carbonyl-containing perfume compounds but must be essentially free of primary amines and ammonia. For purposed hereof, compositions substantially free of primary amines and ammonia includes the

30 protonated as well as unprotonated forms of the compounds.

Once ammonia and primary amines are eliminated from the compositions, it becomes possible to prepare substantially colorless shampoos. However, in order to formulate shampoos which are also mild and high lathering, it is desirable to use certain

35

- 4 -

amphoteric surfactants which typically add some degree of color, usually a yellow tint, to the shampoo. Whereas this is not significant in the formulation of most shampoos (including the various-colored "clear" shampoos) it is undesirable in the formulation of substantially colorless shampoos. Thus, it is also
5 a necessary aspect of this invention that the amphoteric surfactants utilized in the present shampoo compositions be sufficiently colorless and free of primary amines and ammonia such that the shampoo composition itself remains substantially
10 colorless.

The invention, including preferred embodiments thereof, is described in more detail in the Detailed Description of the Invention, which follows.

DETAILED DESCRIPTION OF THE INVENTION

15 The essential as well as a variety of optional components of the compositions of the present invention are described below.

Alkyl Ethoxylated Sulfate

20 The shampoo composition hereof comprises from about 2% to about 20%, by weight, preferably from about 5% to about 15%, more preferably from about 7% to about 15%, of alkyl ethoxylated sulfate anionic surfactant.

25 Alkyl ethoxylated sulfate surfactants are well known in the art, and can be represented by the formula $RO(C_2H_4O)_xSO_3M$, wherein R is alkyl or alkenyl of from about 8 to about 24 carbon atoms, x is 1 to 12, preferably, and M is a water-soluble cation such as an alkali or alkaline earth metal, preferably, sodium or potassium. The average degree of ethoxylated, i.e. the average value for x should be at least about 2.0.

30 Exemplary alkyl ethoxylated sulfates are condensation products of ethylene oxide and monohydric alcohols having from about 8 to about 24 carbon atoms. Preferably, R has from about 10 to about 18 carbon atoms. The alcohols can be derived from fats, e.g., coconut oil or tallow, or can be synthetic. Lauryl alcohol and straight chain alcohols derived from coconut oil are preferred
35 herein. Such alcohols are typically reacted with from about 2 to about 12, preferably about 2 to about 6, more preferably about

- 5 -


about 3, molar proportions of ethylene oxide and the resulting mixture of molecular species having, for example, an average moles of ethylene oxide per mole of alcohol also within the above limits, is sulfated and neutralized.

Specific examples of alkyl ethoxylated sulfates which may be used in the present invention are the salts, especially sodium and/or potassium salts, of coconut alkyl triethylene glycol ethoxylated sulfate, tallow alkyl triethylene glycol ethoxylated sulfate, and tallow alkyl hexaoxyethylene sulfate. Typically the alkyl ether sulfates will comprise a mixture of individual compounds, said mixture preferably having an average alkyl chain length of from about 10 to about 16 carbon atoms, and an average degree of ethoxylation of from about 2 to about 6 moles of ethylene oxide. Especially preferred are narrow range alkyl ethoxylate sulfates such as those having ethoxylation levels primarily in the range of about 2. to about 6.

Amphoteric Surfactant

The amphoteric surfactant will be present in the shampoo compositions hereof at levels of from about 1% to about 15%, by weight of the composition, preferably from about 1.5% to about 10%, more preferably about 2.5% to about 8%. The amphoteric component hereof is selected from the group consisting of amphoteric betaine, imidazoline, aminoalkanoate, and iminodialkanoate surfactants. Preferably, the ratio of the alkyl ethoxylated surfactant to the amphoteric surfactant will be from about 20:1 to about 1:2, more preferably from about 10:1 to about 2:1.

The imidazoline amphoteric surfactants hereof are depicted by Formula I:



wherein R^1 is $\text{C}_8 - \text{C}_{22}$ alkyl or alkenyl, preferably $\text{C}_{12} - \text{C}_{16}$, R^2 is hydrogen or $\text{CH}_2\text{CO}_2\text{M}$, R^3 is $\text{CH}_2\text{CH}_2\text{OH}$ or $\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{COOM}$, R^4 is

- 6 -

hydrogen, $\text{CH}_2\text{CH}_2\text{OH}$, or $\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{COOM}$, Z is CO_2M or $\text{CH}_2\text{CO}_2\text{M}$, n is 2 or 3, preferably 2, M is hydrogen or a cation, such as alkali metal or alkaline earth metal. Examples of "alkali metal" include lithium, sodium, and potassium. Examples of "alkaline earth metal" include beryllium, magnesium, calcium, strontium, and barium. This type of surfactant is classified herein as an "imidazoline" amphoteric surfactant for convenience, although it should be recognized that it does not necessarily have to be derived, directly or indirectly, through an imidazoline intermediate.


Suitable materials of this type are marketed under the tradename MIRANOL and are understood to comprise a complex mixture of species, and can exist in protonated and non-protonated species depending upon pH with respect to species that can have a hydrogen at R^2 . All such variations and species are meant to be encompassed herein.

Preferred surfactants of Formula I are monocarboxylates and dicarboxylates. Examples of these materials include cocoamphocarboxypropionate, cocoamphocarboxypropionic acid, cocoamphocarboxyglycinate (alternately referred to as cocoamphodiacetate), and cocoamphoacetate (alternately, cocoamphomonoacetate).

Specific commercial products providing the amphoteric surfactant component of the present compositions include those sold under the trade names MIRANOL C2M CONC. N.P., MIRANOL C2M CONC. O.P., MIRANOL C2M SF, MIRANOL CM SPECIAL (Miranol, Inc.); ALKATERIC 2CIB (Alkaril Chemicals); AMPHOTERGE W-2 (Lonza, Inc.); MONATERIC CDX-38, MONATERIC CSH-32 (Mona Industries); REWOTERIC AM-2C (Rewo Chemical Group); and SCHERCOTERIC MS-2 (Scher Chemicals).

- 7 -

Suitable betaine surfactants hereof are depicted by compounds having the Formula (II):



wherein:

R_1 is a member selected from the group consisting of
 COOM and $\text{CH}-\text{CH}_2\text{SO}_3\text{M}$
 $\quad \quad \quad |$
 $\quad \quad \quad \text{OH}$

R_2 is lower alkyl or hydroxyalkyl;

R_3 is lower alkyl or hydroxyalkyl;

R_4 is a member selected from the group consisting of hydrogen and lower alkyl;

R_5 is higher alkyl or alkenyl;

Y is lower alkyl, preferably methyl;

m is an integer from 2 to 7, preferably from 2 to 3;

n is the integer 1 or 0;

M is hydrogen or a cation, as previously described, such as an alkali metal or alkaline earth metal.

The term "lower alkyl" or "hydroxyalkyl" means straight or branch chained, saturated, aliphatic hydrocarbon radicals and substituted hydrocarbon radicals having from one to about three carbon atoms such as, for example, methyl, ethyl, propyl, isopropyl, hydroxypropyl, hydroxyethyl, and the like. The term "higher alkyl or alkenyl" means straight or branch chained saturated (i.e., "higher alkyl") and unsaturated (i.e., "higher alkenyl") aliphatic hydrocarbon radicals having from about eight to about 20 carbon atoms such as, for example, lauryl, cetyl, stearyl, oleyl, and the like.


Examples of surfactant betaines of Formula II wherein n is zero which are useful herein include the alkylbetaines such as cocodimethylcarboxymethylbetaine, lauryldimethylcarboxymethylbetaine, lauryl dimethyl- α -carboxyethylbetaine,

- 8 -

cetyldimethylcarboxymethylbetaine,
 lauryl-bis-(2-hydroxyethyl)carboxymethylbetaine,
 stearyl-bis-(2-hydroxypropyl)carboxymethylbetaine, oleyl-
 dimethyl-gamma-carboxypropylbetaine, lauryl-bis-(2-hydroxypropyl)-
 alpha-carboxyethylbetaine, etc. The sulfobetaines may be repre-
 5 sented by cocodimethylsulfopropylbetaine, stearyldimethylsulfo-
 propylbetaine, lauryl-bis-(2-hydroxyethyl)sulfopropylbetaine, and
 the like.

Amido betaines and amidosulfo betaine surfactants useful in
 the present invention are exemplified by compounds of Formula II
 10 wherein n is one but otherwise corresponding to the above ex-
 amples. Examples of surfactant betaines of Formula II wherein n
 is one which are useful herein include the amidocarboxybetaines,
 such as cocoamidodimethylcarboxymethylbetaine, laurylamidodi-
 methylcarboxymethylbetaine, cetylamidodimethylcarboxymethyl-
 15 betaine, laurylamido-bis-(2-hydroxyethyl)-carboxymethylbetaine,
 cocoamido-bis-(2-hydroxyethyl)-carboxymethylbetaine, etc. The
 amido sulfobetaines may be represented by cocoamidodimethylsulfo-
 propylbetaine, stearylamidodimethylsulfopropylbetaine, lauryl-
 amido-bis-(2-hydroxyethyl)-sulfopropylbetaine, and the like.
 20

The preferred betaine in the present invention is a member
 selected from the group consisting of surfactant amidocarboxy-
 betaines and amidosulfobetaines. More preferred betaines are the
 surfactant amidocarboxybetaines, particularly cocoamidodimethyl-
 carboxymethylbetaines (cocamidopropylbetaine), such as those sold
 25 by Goldschmidt Co. under the trade name Tegobetaine L-7(R grade),
 and by Hoechst-Celanese under the trade name Genagen CAB. These
 most preferred betaines have the formula:



wherein R'₃ is selected from C8 to C18 alkyl radicals and M is
 hydrogen or a cation as defined above. In general, the preferred

35

- 9 -

betaines hereof will have low levels of residual amide and sodium monochloroacetate.

Suitable aminoalkanoates and iminodialkanoates are represented by the Formulas (III) and (IV):

5 aminoalkanoates of the formula:



10 iminodialkanoates of the formula:



15 wherein n and m are from 1 to 4, R is C₈ - C₂₂ alkyl or alkenyl, and M is hydrogen or alkali or an alkaline earth metal as previously described.

20 Examples of amphoteric surfactants falling within the aminoalkanoate formula include n-alkylamino-propionates and n-alkyliminodipropionates. Such material are sold under the tradename DERIPHAT by Henkel and MIRATANE by Miranol, Inc. Specific examples include N-lauryl-beta-amino propionic acid or salts thereof, and N-lauryl-beta-imino-dipropionic acid (DERIPHAT 160C) or salts thereof, and mixtures thereof.

25 The amphoteric surfactant provided as a raw material for use in the shampoo compositions hereof must be sufficiently colorless such that it does not impact enough color to the total composition remains substantially colorless.

30 Amphoteric surfactants are often used in conventional shampoo compositions at a level and supplied in a form that tend to impart a degree of color in excess of that which is preferred for the compositions herein. The amphoteric surfactants utilized in the present compositions must be sufficiently colorless so that the overall absorbance value of the final product is within the limits set forth herein. Reduced color amphoteric surfactants raw materials can easily be produced by those skilled in the art, for

35

- 10 -

instance, by the use of cleaner, purer raw materials and minimizing the period of time the surfactant or reaction mixture is exposed to elevated temperature during manufacture of the surfactant. Suitable surfactants are currently commercially available. e. g. Tegobetaine L-7 (R Grade) from Goldschmidt.

5 Preferably, the amphoteric surfactant, as well as the other surfactants added to the composition, will exhibit an absorbance value in a 30% aqueous solution at 440nm in a 1.0cm path length of no greater than about 0.1, more preferably no greater than about 0.05, most preferably no greater than about 0.035.

10 N-Acylamino Acid Surfactant

The shampoo compositions of the present invention comprise from about 0.5% to about 10%, preferable from about 0.5% to about 5%, more preferably from about 1% to about 5%, of N-acyl amino acid surfactant.

15 N-acyl amino acid surfactants, for purposes hereof, include N-acyl hydrocarbyl acids and salts thereof, such as those represented by Formula V, as follows:



20 wherein: R¹ is a C₈-C₂₄ alkyl or alkenyl radical, preferably C₁₂-C₁₈; R² is -H, C₁-C₄ alkyl, phenyl, or -CH₂COOM, preferably C₁-C₄ alkyl, more preferably C₁-C₂ alkyl; R³ is -CR⁴₂- or C₁-C₂ alkoxy, wherein each R⁴ independently is -H or C₁-C₆ alkyl or alkylester, and n is from 1 to 4, preferably 1 or 2; and M is -H or a cation as previously defined, preferably an alkali metal such as sodium or potassium.

30 A wide variety of N-acyl acid surfactants and their synthesis are described in Anionic Surfactants, Part II, Surfactant Science Series, Vol. VII, edited by Warner M. Linfield, Marcel Dekker, Inc. (New York and Basel), 1976; pp 581-617.

Especially preferred are compounds of Formula V wherein R² is methyl and R³ is -CH₂-, and n is 1, which are known as the N-acyl sarcosinates, and acids thereof. Specific examples include

35

- 11 -

lauroyl sarcosinate, myristoyl sarcosinate, cocoyl sarcosinate, and oleoyl sarcosinate, preferably in their sodium and potassium salt forms.

5 For the purposes of the surfactants described herein, it should be understood that the terms "alkyl" or "alkenyl" include mixtures of radicals which may contain one or more intermediate linkages such as ether or polyether linkages or non-functional substituents such as hydroxyl or halogen radicals wherein the radical remains of hydrophobic character.

10 Perfume

Perfumes are a highly desirable element of hair care compositions, since they can mask undesirable odors of other ingredients of the shampoo, as well as provide a pleasing aesthetic fragrance. Accordingly, perfume is an essential ingredient in the present compositions.

15 A wide variety of perfumes are known to those skilled in the art and are commercially available. The particular perfume used is largely a matter of choice. However the perfume should be used at a level effective for providing a noticeable aroma to the composition, or for masking undesired aroma of the composition.

20 In general, the compositions will comprise from about 0.05% to about 10% of a perfume compounds containing carbonyl functionalities, preferably from about 0.1% to about 7%, more preferably from about 0.1% to about 3%. As used herein, perfume compound means aromatically active ingredients whereas "perfume" or "perfume component" includes the total perfume compounds and any accompanying perfume solvent.

25 Perfumes are made by those skilled in the art in a wide variety of fragrances and strengths. Typical perfumes are described in Arctander, Perfume and Flavour Chemicals (Aroma Chemicals), Vol. I and II (1969); and Arctander, Perfume and Flavour Materials of Natural Origin (1960).

30 Carbonyl-containing perfume compounds include aldehydes, ketones and esters. These materials can be aliphatic (saturated

- 12 -

or unsaturated), aromatic, heterocyclic, or alicyclic. Generally they will have from about 6 to about 18 carbon atoms.

Aliphatic aldehyde perfume compounds include hexyl aldehyde, heptyl aldehyde, octyl aldehyde, decyl aldehyde, octyl aldehyde, decyl aldehyde, undecyl aldehyde, undecylinic aldehyde, lauric aldehyde, methyl monyl acetaldehyde, myristic aldehyde, and nonyl aldehyde. Aromatic aldehydes include cinnamic aldehydes such as amyl cinnamic aldehyde and hexyl cinnamic aldehyde, anisic aldehyde, heliotropin, benzaldehyde, and vanillin. Heterocyclic aldehydes include piperonal and furan.

Aliphatic ketone perfume compounds include hexanone, heptanone, octanone, ionone, undecalactone, nonalactone (gamma-nonyl lactone), camphor, alpha-methyl ionone, (gamma-undecyl lactone) gamma-methyl ionone, and beta-methyl naphthyl ketone. Alicyclic ketones include, for example, amyl cyclohexanone. Aromatic ketones include beta-naphthyl methyl ketones, para-hydroxy phenolbutanone, laevo carvone, and musk ketone. Heterocyclic ketones include thujone, cedrenone, and exaltolide.

Aliphatic ester perfume compounds include acetates, butyrates, and formates, such as ethyl butyrate, ethyl formate, butyl acetate, isopropyl butyrate, ethyl formate, cis-3-hexenyl acetate, ethyl aceto acetate, and phenylethyl phenylacetate, carbonates such as methyl octine carbonate, and phthalates such as diethyl phthalate. Heterocyclic esters include cedryl formate and cedryl acetate. Aromatic esters include benzyl acetate, benzyl salicylate, cis-3-hexenyl acetate, cis-3-hexenyl salicylate, and methyl anthranilate. Other esters include alicyclic esters and aliphatic/alicyclic esters, such as 4 - tertiary butyl cyclohexyl acetate. Ester perfumes also include materials derived from floral materials and fruits, and from animal notes (e.g., natural isolates of civet, castoreum, and musk).

Perfume solvents are well known in the art, and the conventional ones can be used herein, e.g., dipropylene glycol, diethylene glycol, C₁-C₆ alcohols, etc.

- 13 -

Water

The shampoo composition hereof will also comprise water. Generally, the composition will contain from about 50% to about 90% water, preferably about 60% to about 85% water, more preferably from about 65% to about 85%.

Shampoo Compositions

The shampoo hereof is substantially free of alkyl sulfate surfactants since alkyl sulfates are relatively harsh to the skin. It is recognized that there will generally be some alkyl sulfate present as a result of it being present in commercially available alkyl ethoxylated sulfate raw materials. For example, commercially available alkyl (3) ethoxylated sulfate typically contains about 20% by weight alkyl sulfate; commercially available alkyl (2) ethoxylated sulfate, about 25% to about 40% alkyl sulfate. For purposes hereof, substantially free of alkyl sulfate means the compositions hereof should have an alkyl sulfate:alkyl ethoxylated sulfate (average degree of ethoxylation of 2.5 and above) weight ratio of no more than about 0.35, preferably no more than about 0.30, more preferably no more than about 0.25. For alkyl ethoxylated sulfate with an average ethoxylation level of less than 2.5, the ratio should be no more than about 0.40, preferably no more than about 0.35, more preferably no more than about 0.30, most preferably no more than about 0.25. It is preferred that no additional amount of alkyl sulfate be added other than that which occurs inherently with the alkyl ethoxylated sulfate.

Narrow range ethoxylates can be used to lower the alkyl sulfate: alkyl ethoxylated sulfate weight ratio. "Narrow range ethoxylates" refer to alkyl ethoxylated sulfate surfactants that have been processed to reduce alkyl sulfates and, optionally, alkyl ethoxylated sulfates outside of the desired range of ethoxylation. The use of narrow range ethoxylates can be used to lower the alkyl sulfate:alkyl ethoxylated sulfate weight ratio, including to ratios as low as about 0.2 or even about 0.1, and less.

- 14 -

The present shampoo compositions are preferably substantially free of amide foam boosters such as fatty (e.g. C₁₂-C₂₀) mono- and di- alkanol amides, since these materials can be harsh to the skin and are not needed in the present compositions. "Substantially free of amide foam boosters" means the compositions hereof can contain no more than about 1.0%, by weight of the amide foam booster, preferably no more than about 0.5%, more preferably no more than about 0.2%. Most preferably, no amide foam booster is added.

As an essential aspect hereof, the shampoo compositions are substantially free of ammonia and primary amines. "Substantially free of ammonium ions and free primary amines" means that the compositions hereof can contain no more than about 0.1%, by weight, of ammonia (including ammonium ions) and primary amines (including free amines and protonated primary amines), preferably no more than about 0.05%.

It is also preferred that no other ingredients that are unduly harsh to the skin be added to the shampoo compositions hereof.

The shampoo compositions of the present invention are substantially colorless, and can remain substantially colorless over prolonged periods of time. Substantially colorless, as used herein, means the shampoos are both clear and are characterized by lack of color. More specifically, the shampoo compositions hereof have an absorbance value at 440nm in a 1.0cm path length of no greater than about 0.030 after storage at 38°C for 30 days, preferably no greater than about 0.025. Absorbance value is measured utilizing a spectrophotometer in accordance with conventional techniques known in the art.

Additional Ingredients

The compositions of the present invention can contain a wide variety of optional ingredients useful or known for use in the art for shampoo and other hair and skin cleansing compositions. Exemplary additional ingredients are described below.

Additional surfactants that can be used include other


- 15 -

anionic, nonionic, and amphoteric surfactants, as well as zwitter-ionic and cationic surfactants.

Anionic Surfactants

A suitable class of anionic surfactants are the water-soluble, organic salts of the general formula:

5



wherein R_1 is chosen from the group consisting of a straight or branched chain, saturated aliphatic hydrocarbon radical having from about 8 to about 24, preferably about 12 to about 18, carbon atoms; and M is a cation. Important examples are the salts of an organic sulfuric acid reaction product of a hydrocarbon of the methane series, including iso-, neo-, and n-paraffins, having about 8 to about 24 carbon atoms, preferably about 12 to about 18 carbon atoms and a sulfonating agent, e.g., SO_3 , H_2SO_4 , oleum, obtained according to known sulfonation methods, including bleaching and hydrolysis. Preferred are alkali metal sulfonated C₁₂₋₁₈ paraffins.

10

Additional examples of anionic surfactants which come within the terms of the present invention are the reaction products of fatty acids esterified with isethionic acid and neutralized with sodium hydroxide where, for example, the fatty acids are derived from coconut oil; sodium or potassium salts of fatty acid amides of methyl tauride in which the fatty acids, for example, are derived from coconut oil. Other anionic synthetic surfactants of this variety are set forth in U.S. Patents 2,486,921; 2,486,922; and 2,396,278.

20

Still other anionic surfactants include the class designated as succinamates. This class includes such surface active agents as disodium N-octadecylsulfosuccinamate; tetrasodium N-(1,2-dicarboxyethyl)-N-octadecylsulfosuccinamate; diamyl ester of sodium sulfosuccinic acid; dihexyl ester of sodium sulfosuccinic acid; dioctyl esters of sodium sulfosuccinic acid.

25

Other suitable anionic surfactants utilizable herein are olefin sulfonates having about 12 to about 24 carbon atoms. The term "olefin sulfonates" is used herein to mean compounds which


30

35

- 16 -

can be produced by the sulfonation of α -olefins by means of uncomplexed sulfur trioxide, followed by neutralization of the acid reaction mixture in conditions such that any sulfones which have been formed in the reaction are hydrolyzed to give the corresponding hydroxy-alkanesulfonates. The sulfur trioxide can be liquid or gaseous, and is usually, but not necessarily, diluted by inert diluents, for example by liquid SO₂, chlorinated hydrocarbons, etc., when used in the liquid form, or by air, nitrogen, gaseous SO₂, etc., when used in the gaseous form.

Another class of anionic surfactants are the β -alkyloxy alkane sulfonates. These compounds have the following formula:



where R₁ is a straight chain alkyl group having from about 6 to about 20 carbon atoms, R₂ is a lower alkyl group having from about 1 (preferred) to about 3 carbon atoms, and M is a water-soluble cation.

Many additional synthetic anionic surfactants are described in McCutcheon's, Emulsifiers and Detergents, 1989 Annual, published by M. C. Publishing Co., which is incorporated herein by reference. Also U.S. Patent 3,929,678, Laughlin et al., issued December 30, 1975, discloses many other anionic as well as other surfactant types and is incorporated herein by reference. Soaps, of course, also fall within the scope of anionic deterative surfactants that can be used.

Nonionic Surfactants

A wide variety of nonionic surfactants can be used. Nonionic surfactants include those broadly defined as compounds produced by the condensation of alkylene oxide groups (hydrophilic in nature) with an organic hydrophobic compound, which may be aliphatic or

- 17 -

alkyl aromatic in nature. Examples of classes of nonionic surfactants are:

5 1. The polyethylene oxide condensates of alkyl phenols, e.g., the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 20 carbon atoms, preferably from about 6 to about 12, in either a straight chain or branched chain configuration, with ethylene oxide, the said ethylene oxide being present in amounts equal to from about 10 to about 60 moles of ethylene oxide per mole of alkyl phenol. The alkyl substituent
10 in such compounds may be derived from polymerized propylene, diisobutylene, octane, or nonane, for example.

2. Those derived from the condensation of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylene diamine products which may be varied in composition
15 depending upon the balance between the hydrophobic and hydrophilic elements which is desired. For example, compounds containing from about 40% to about 80% polyoxyethylene by weight and having a molecular weight of from about 5,000 to about 11,000 resulting from the reaction of ethylene oxide groups with a hydrophobic base constituted of the reaction product of ethylene diamine and excess
20 propylene oxide, said base having a molecular weight of the order of about 2,500 to about 3,000, are satisfactory.

3. The condensation product of aliphatic alcohols having from about 8 to about 18 carbon atoms, in either straight chain or branched chain configuration, with ethylene oxide, e.g., a coconut
25 alcohol ethylene oxide condensate having from about 10 to about 30 moles of ethylene oxide per mole of coconut alcohol, the coconut alcohol fraction having from about 10 to about 14 carbon atoms.

4. Long chain tertiary amine oxides corresponding to the following general formula:

30
$$R_1R_2R_3N \text{ -----} \rightarrow O$$

wherein R_1 contains an alkyl, alkenyl or monohydroxy alkyl radical of from about 8 to about 18 carbon atoms, from 0 to about 10 ethylene oxide moieties, and from 0 to about 1 glyceryl moiety, and R_2 and R_3 contain from about 1 to about 3 carbon atoms and
35

- 18 -

from 0 to about 1 hydroxy group, e.g., methyl, ethyl, propyl, hydroxyethyl, or hydroxypropyl radicals. The arrow in the formula is a conventional representation of a semipolar bond. Examples of amine oxides suitable for use in this invention include dimethyl-
 5 dodecylamine oxide, oleyldi(2-hydroxyethyl) amine oxide, dimethyl-octylamine oxide, dimethyl-decylamine oxide, dimethyl-tetradecyl-amine oxide, 3,6,9-trioxaheptadecyldiethylamine oxide, di(2-hydroxyethyl)-tetradecylamine oxide, 2-dodecoxyethyl dimethylamine oxide, 3-dodecoxy-2-hydroxypropyldi(3-hydroxypropyl) amine oxide,
 10 dimethylhexadecylamine oxide.

5. Long chain tertiary phosphine oxides corresponding to the following general formula:



wherein R contains an alkyl, alkenyl or monohydroxyalkyl radical ranging from about 8 to about 18 carbon atoms in chain length,
 15 from 0 to about 10 ethylene oxide moieties and from 0 to about 1 glyceryl moiety and R' and R'' are each alkyl or monohydroxyalkyl groups containing from about 1 to about 3 carbon atoms. The arrow in the formula is a conventional representation of a semipolar bond. Examples of suitable phosphine oxides are:
 20 dodecyldimethylphosphine oxide, tetradecyldimethylphosphine oxide, tetradecylmethylethylphosphine oxide, 3,6,9,-trioxaoctadecyldimethylphosphine oxide, cetyldimethylphosphine oxide, 3-dodecoxy-2-hydroxypropyldi(2-hydroxyethyl) phosphine oxide, stearyldimethyl-
 25 phosphine oxide, cetylethylpropylphosphine oxide, oleyldiethylphosphine oxide, dodecyldiethylphosphine oxide, tetradecyldiethylphosphine oxide, dodecyldipropylphosphine oxide, dodecyldi(hydroxymethyl)phosphine oxide, dodecyldi(2-hydroxyethyl)phosphine oxide, tetradecylmethyl-2-hydroxypropylphosphine oxide, oleydimethylphosphine oxide, 2-hydroxydodecyldimethylphosphine oxide.

30 6. Long chain dialkyl sulfoxides containing one short chain alkyl or hydroxy alkyl radical of from about 1 to about 3 carbon atoms (usually methyl) and one long hydrophobic chain which include alkyl, alkenyl, hydroxy alkyl, or keto alkyl radicals
 35 containing from about 8 to about 20 carbon atoms, from 0 to about

10 ethylene oxide moieties and from 0 to about 1 glyceryl moiety. Examples include: octadecyl methyl sulfoxide, 2-ketotridecyl methyl sulfoxide, 3,6,9,-trioctadecyl 2-hydroxyethyl sulfoxide, dodecyl methyl sulfoxide, oleyl 3-hydroxypropyl sulfoxide, tetradecyl methyl sulfoxide, 3-methoxytridecyl methyl sulfoxide, 3-hydroxytridecyl methyl sulfoxide, 3-hydroxy-4-dodecoxybutyl methyl sulfoxide.

5
10 7. Polysorbates, e.g., sucrose esters of fatty acids. Such materials are described in U.S. Patent 3,480,616, e.g., sucrose cocoate (a mixture of sucrose esters of a coconut acid, consisting primarily of monoesters, and sold under the tradenames GRILLOTEN LSE 87K from RITA, and CRODESTA SL-40 from Croda).

15 8. Alkyl polysaccharide nonionic surfactants are disclosed in U.S. Patent 4,565,647, Llenado, issued January 21, 1986, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g., a polyglycoside, hydrophilic group. The polysaccharide can contain from about 1.0 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties. (Optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside.) The intersaccharide bonds can be, 25 e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-positions on the preceding saccharide units.

30 Optionally, and less desirably, there can be a polyalkylene-oxide chain joining the hydrophobic moiety and the polysaccharide moiety. The preferred alkyleneoxide is ethylene oxide. Typical hydrophobic groups include alkyl groups, either saturated or unsaturated, branched or unbranched containing from about 8 to about 18, preferably from about 10 to about 16, carbon atoms. 35 Preferably, the alkyl group is a straight chain saturated alkyl

group. The alkyl group can contain up to about 3 hydroxy groups and/or the polyalkyleneoxide chain can contain up to about 10, preferably less than 5, alkylene moieties. Suitable alkyl polysaccharides are octyl, nonyldecyl, undecyldodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, and octadecyl, di-, tri-, tetra-, penta-, and hexagluco-
sides, galactosides, lactosides, glucoses, fructosides, fructoses and/or galactoses. Suitable mixtures include coconut alkyl, di-, tri-, tetra-, and pentagluco-
sides and tallow alkyl, tetra-, penta-, and hexagluco-
sides.

9. Polyethylene glycol (PEG) glyceryl fatty esters, as depicted by the formula $RC(O)OCH_2CH(OH)CH_2(OCH_2CH_2)_nOH$ wherein n is from about 5 to about 200, preferably from about 20 to about 100, more preferably from about 30 to about 85, and $RC(O)-$ is an ester wherein R comprises an aliphatic radical having from about 7 to 19 carbon atoms, preferably from about 9 to 17 carbon atoms, more preferably from about 11 to 17 carbon atoms, most preferably from about 11 to 14 carbon atoms. The combinations of n from about 20 to about 100, with C_{12} - C_{18} , preferably C_{12} - C_{15} fatty esters, for minimized adverse effect on foaming, is preferred.

Suitable glyceryl fatty ester portions of these surfactants include glyceryl cocoate, glyceryl tallowate, glyceryl palmate, glyceryl stearate, glyceryl laurate, glyceryl oleate, glyceryl ricinoleate, and glyceryl fatty esters derived from triglycerides, such as palm oil, almond oil, and corn oil.

Other surfactants that can be used include soluble cationic surfactants, such as quaternary ammonium surfactants, and other amphoteric and zwitterionic surfactants known to those in the art.

Conditioning Agent

An optional component hereof is a soluble conditioning agent suitable for conditioning hair or skin.

Suitable conditioners include, for example, soluble polyether siloxane copolymer, such as a polypropylene oxide modified dimethylpolysiloxane (e.g., Dow Corning DC-1248), although ethylene oxide or mixtures of ethylene oxide and

- 21 -

propylene oxide may also be used. The ethylene oxide and polypropylene oxide level must be sufficiently high to provide solubility in water and the composition hereof.

Antidandruff Agent

5 The shampoo compositions can also contain one or more antidandruff agents. Soluble antidandruff agents should be used.

Hydroxypyridone salts are disclosed in *Cosmetics and Drug Preservation, Principles and Practice*, p 742 (edited by J. Kabura, 1984), incorporated herein by reference. Hydroxypyridone salts used herein include those generally described as 1-hydroxy-4-methyl-(1H)-pyridones having an aliphatic or aromatic moiety (R) at the 6 position thereof, wherein R has a factor of at least 1:3, preferably from 2 to 6, more preferably from 3 to 5.5. The factor is a measure of the lipophilicity/ hydrophilicity of the substituent and is defined in detail in the paper by W. Dittmar, E. Druckrey and H. Urbach, *J. Med. Chem.* 17(7), 753-6 (1974) and references cited therein; all of which are incorporated herein by reference.

10 In structural terms, preferred R substituents are selected from linear and branched C₃-C₁₁, preferably C₄-C₁₁, alkyl and alkenyl groups, C₃-C₈ cycloalkyl groups, and C₃-C₈, preferably C₆, aryl groups. The cyclic moieties, discussed above, can also be substituted with one or more alkyl or alkenyl groups up to C₄. The R groups can further be substituted with halogen atoms. Of the above, preferred R moieties are cyclohexyl and 2,4,4-trimethyl pentyl, the latter being highly preferred.

25 The above mentioned compounds can be used both in the free form and as salts, for example, salts with organic bases or inorganic cations. Low molecular weight alkanolamines are especially preferred organic bases. The preferred hydroxypyridone salt for use herein is monoethanolamine salt known as piroctone olamine or Octopirox; see *Cosmetic and Drug Preservation, supra*.

Other Optional Components

35 The compositions herein can contain a variety of other non-essential optional components. Such optional ingredients

- 22 -

include, for example, preservatives such as benzyl alcohol, methyl paraben, propyl paraben and imidazolidinyl urea; cationic conditioning agents, including both cationic conditioning surfactant and cationic conditioning polymers; fatty alcohols; block polymers of ethylene oxide and propylene oxide such as Pluronic F88 offered by BASF Wyandotte; viscosity modifiers, such as sodium chloride, sodium sulfate, and xylene sulfonate salts; propylene glycol; polyvinyl alcohol; ethyl alcohol; foam boosters such as Polyquaternium-10 (an industry term designated by The Cosmetic, Toiletry and Fragrance Association (CTFA) for the polymeric quaternary ammonium salt of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide), commercially available from Union Carbide Corp. (Danbury, Connecticut, USA) under their UCARE POLYMER JR series of materials, e.g., UCARE POLYMER JR-30M, JR-125 and JR-400; additional pH adjusting agents for the final shampoo composition such as citric acid, succinic acid, phosphoric acid, sodium hydroxide, sodium carbonate, etc.; perfumes; dyes; and sequestering agents such as disodium ethylene-diamine tetraacetate.

Another type of suspending agent that can be used is carboxyvinyl polymer. Preferred polymers are copolymers of acrylic acid crosslinked with polyallylsucrose as described in U.S. Patent 2,798,053, Brown, issued July 2, 1957, incorporated herein by reference. These polymers are provided by B. F. Goodrich Company as, for example, Carbopol 934, 940, 941, and 956.

Other materials can also be used as suspension agents, including those that can impart a gel-like viscosity to the composition, such as water soluble or colloiddally water soluble polymers like cellulose ethers (e.g., hydroxyethyl cellulose), guar gum, polyvinyl alcohol, polyvinyl pyrrolidone, hydroxypropyl guar gum, starch and starch derivatives, and other thickeners, viscosity modifiers, gelling agents, etc. Mixtures of these materials can also be used.

Another type of suspending agent that can be used is xanthan gum. Shampoo compositions utilizing xanthan gum as a suspending

- 23 -

agent for the silicone hair conditioning component are described in U.S. Patent 4,788,006, Bolich and Williams, issued November 29, 1988, incorporated herein by reference. Xanthan gum is bio-synthetic gum material that is commercially available.

5 Optional ingredients are typically used at levels of from about 0.01% to about 10% of the composition. This list of optional ingredients is not meant to be inclusive, as other optional components can be utilized.

10 The pH of the compositions for direct use for cleaning applications is preferably in the range of from 2 to about 10, more preferably from about 4 to about 8, most preferably from about 5 to about 8.

METHOD OF USE

15 The present compositions are used in a conventional manner for cleaning hair. An effective amount of the composition for cleaning hair, typically, from about 1 g to about 20 g of the composition, is applied to hair that has preferably been wetted, generally with water, and then rinsed out. Application to the hair typically includes working the composition through the hair
20 such that most or all of the hair is contacted with the composition.

EXAMPLES

The following Examples further describe and demonstrate the preferred embodiments within the scope of the present invention.
25 The Examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention as many variations thereof are possible without departing from its spirit and scope.

30

35

- 24 -

EXAMPLES I-XX

The following examples exemplify shampoo compositions of the present invention.

<u>Component (active wt.%)</u>		<u>I</u>	<u>II</u>	<u>III</u>	<u>IV</u>	<u>V</u>
5	Sodium Laureth-3 Sulfate (%) ¹	10	13.5	10	13.5	13.5
	Cocoamidopropyl Betaine (%) ²	5	1.5	5	1.5	3.5
	Sodium Lauroyl Sarcosinate (%) ³	4.5	4.5	1.5	1.5	3.5
	Cocoamphodiacetate (%) ⁴	-	-	-	-	-
	Lauroiminodiacetate (%) ⁵	-	-	-	-	-
10	Lauroiminodipropionate (%) ⁶	-	-	-	-	-
	Dimethicone Copolyol (%) ⁷	-	-	-	-	-
	Polyol alkoxyester (%) ⁸	-	-	-	-	-
	Piroctone olamine (%)	-	-	-	-	-
	Perfume (%) ⁹	0.3	0.3	0.3	0.3	0.3
15	DMDM Hydantoin (%)	0.37	0.37	0.37	0.37	0.37
	EDTA (%)	0.1	0.1	0.1	0.1	0.1
	Citric Acid	0.2	0.2	0.2	0.2	0.2
	NaCl	2	4	1	1	1
	Water	q.s.	q.s.	q.s.	q.s.	q.s.
<u>Component (active wt.%)</u>		<u>VI</u>	<u>VII</u>	<u>VIII</u>	<u>IX</u>	<u>X</u>
20	Sodium Laureth-3 Sulfate (%) ¹	13	12.8	13.5	13.5	13.5
	Cocoamidopropyl Betaine (%) ²	5	6	-	-	-
	Sodium Lauroyl Sarcosinate (%) ³	1	2	3.5	3.5	3.5
	Cocoamphodiacetate (%) ⁴	-	-	3.5	-	-
	Lauroiminodiacetate (%) ⁵	-	-	-	3.5	-
25	Lauroiminodipropionate (%) ⁶	-	-	-	-	3.5
	Dimethicone Copolyol (%) ⁷	-	-	-	-	-
	Polyol alkoxyester (%) ⁸	-	-	-	-	-
	Piroctone olamine (%)	-	-	-	-	-
	Perfume (%) ⁹	0.3	0.3	0.3	0.3	0.3
30	DMDM Hydantoin (%)	0.37	0.37	0.37	0.37	0.37
	EDTA (%)	0.1	0.1	0.1	0.1	0.1
	Citric Acid	0.2	0.2	-	-	-
	NaCl	1	1	1	1	1
	Water	q.s.	q.s.	q.s.	q.s.	q.s.
35						

- 25 -

<u>Component (active wt.%)</u>		<u>XI</u>	<u>XII</u>	<u>XIII</u>	<u>XIV</u>	<u>XV</u>
	Sodium Laureth-3 Sulfate (%) ¹	10	10	10	13.5	13.5
	Cocoamidopropyl Betaine (%) ²	5	5	5	-	-
	Sodium Lauroyl Sarcosinate (%) ³	4.5	4.5	4.5	4.5	4.5
5	Cocoamphodiacetate (%) ⁴	-	-	-	1.5	-
	Lauroiminodiacetate (%) ⁵	-	-	-	-	1.5
	Lauroiminodipropionate (%) ⁶	-	-	-	-	-
	Dimethicone Copolyol (%) ⁷	0.5	-	-	-	-
	Polyol alkoxyester (%) ⁸	-	1.5	-	-	-
10	Piroctone olamine (%)	-	-	1.0	-	-
	Perfume (%) ⁹	0.3	0.3	0.3	0.3	0.3
	DMDM Hydantoin (%)	0.37	0.37	0.37	0.37	0.37
	EDTA (%)	0.1	0.1	0.1	0.1	0.1
	Citric Acid	0.2	0.2	0.2	-	-
15	NaCl	2	-	1	1	1
	Water	q.s.	q.s.	q.s.	q.s.	q.s.
<u>Component (active wt.%)</u>		<u>XVI</u>	<u>XVII</u>	<u>XVIII</u>	<u>XIX</u>	<u>XX</u>
	Sodium Laureth-3 Sulfate (%) ¹	13.5	11	13.5	11	10
	Cocoamidopropyl Betaine (%) ²	-	5.5	1.5	5.5	5
20	Sodium Lauroyl Sarcosinate (%) ³	4.5	1.6	4.5	1.6	0.5
	Cocoamphodiacetate (%) ⁴	-	-	-	-	-
	Lauroiminodiacetate (%) ⁵	-	-	-	-	-
	Lauroiminodipropionate (%) ⁶	1.5	-	-	-	-
	Dimethicone Copolyol (%) ⁷	-	-	-	-	-
25	Polyol alkoxyester (%) ⁸	-	-	1.5	-	-
	Piroctone olamine (%)	-	-	-	1.0	-
	Perfume (%) ⁹	0.3	0.3	0.3	0.3	0.3
	DMDM Hydantoin (%)	0.37	0.37	0.37	0.37	0.37
	EDTA (%)	0.1	0.1	0.1	0.1	0.1
30	Citric Acid	-	0.2	-	-	0.2
	NaCl	1	1	-	1	1
	Water	q.s.	q.s.	q.s.	q.s.	q.s.

1 Absorbance value of no greater than about 0.025.

2 Absorbance value of no greater than about 0.035.

35

- 26 -

- 3 Absorbance value of no greater than about 0.025.
- 4 Absorbance value of no greater than about 0.035.
- 5 Absorbance value of no greater than about 0.035.
- 6 Absorbance value of no greater than about 0.035.
- 7 DOW 190, from Dow Chemical USA (Midland, Michigan).
- 5 8 CROTHIX, from Croda, Inc. (New York, New York).
- 9 Total perfume weight percent, containing about 75%, by weight of the total perfume, of carbonyl-containing perfume compounds.

10 The compositions are prepared as follows. The ethoxylated sulfate, sarcosinate, and amphoteric surfactant are mixed together in a tank and heated to 55°C. The EDTA is then dissolved in the mixture. As applicable, the piroctone olamine, dimethicone copolyol, and polyol alkoxyester are added to the composition with stirring. The perfume and DMDM Hydantoin are added and stirred in next. Finally the NaCl and citric acid are added and the mixture is stirred.

15 The compositions of the Examples can provide excellent in-use hair cleaning, lather, mildness and are substantially colorless after 30 days at 38°C.

20

25

30

35

CLAIMS:


1. A mild, substantially colorless shampoo composition characterized in that it comprises:

- (a) from 2% to 20%, by weight, of an alkyl ethoxylated sulfate anionic surfactant having an average degree of ethoxylation of at least 2.0;
- (b) from 1% to 15% of an amphoteric surfactant selected from the group consisting of betaine surfactants, imidazoline surfactants, aminoalkanoate surfactants, and iminodialkanoate surfactants, and mixtures thereof;
- (c) from 0.5% to 10%, by weight of an N-acylamino acid surfactant, or salt thereof;
- (d) at least 0.05%, by weight of the composition, of carbonyl-containing perfume compounds; and
- (e) water;


wherein said shampoo composition is substantially free of alkyl sulfate anionic surfactant, primary amines, and ammonia, and said shampoo composition has an absorbance value at 440nm in a 1.0cm path length of no greater than 0.030 after storage at 38°C for 30 days.

2. A shampoo composition according to Claim 1, wherein said amphoteric surfactant is selected from the group consisting of

(i) betaine surfactants of the formula:



wherein R_1 is a member selected from the group consisting of



R_2 is lower alkyl or hydroxyalkyl;
 R_3 is lower alkyl or hydroxyalkyl;

- 28 -

R_4 is a member selected from the group consisting of hydrogen and lower alkyl;

R_5 is higher alkyl or alkenyl;


Y is lower alkyl, preferably methyl;

m is an integer from 2 to 7;

n is the integer 1 or 0;

M is hydrogen or a cation;

(ii) amphoteric surfactant having the formula




wherein R^1 is C_8 - C_{22} alkyl or alkenyl, R^2 is hydrogen or CH_2CO_2M , R^3 is CH_2CH_2OH or $CH_2CHOCH_2CH_2COOM$, R^4 is hydrogen, CH_2CH_2OH , or $CH_2CH_2OCH_2CH_2COOM$, Z is CO_2M or CH_2CO_2M , n is 2 or 3, and M is hydrogen or a cation; and

(iii) aminoalkanoates of the formula:




iminodialkanoates of the formula:



wherein n and m are from 1 to 4, R is C_8 - C_{22} alkyl or alkenyl, and M is hydrogen or alkali or alkaline with metal as previously described.

(iv) and mixtures thereof.

3. A shampoo composition as in Claim 1 or 2, wherein said N-acyl amino acid surfactant, or salt thereof, is of the formula:



- 29 -

R_4 is a member selected from the group consisting of hydrogen and lower alkyl;

R_5 is higher alkyl or alkenyl;

Y is lower alkyl, preferably methyl;

5 m is an integer from 2 to 7;

n is the integer 1 or 0;

M is hydrogen or a cation;

(ii) 2% to 10%, by weight, of an amphoteric surfactant having the formula



15 wherein R^1 is C_8 - C_{22} alkyl or alkenyl, R^2 is hydrogen or CH_2CO_2M , R^3 is CH_2CH_2OH or $CH_2CHOCH_2CH_2COOM$, R^4 is hydrogen, CH_2CH_2OH , or $CH_2CH_2OCH_2CH_2COOM$, Z is CO_2M or CH_2CO_2M , n is 2 or 3, and M is hydrogen or a cation; and

(iii) aminoalkanoates of the formula:




iminodialkanoates of the formula:



wherein n and m are from 1 to 4, R is C_8 - C_{22} alkyl or alkenyl, and M is hydrogen or alkali or alkaline with metal as previously described.

30 (iv) and mixtures thereof.

3. A shampoo composition as in Claim 2, wherein said N-acyl amino acid surfactant, or salt thereof, is of the formula:



- 30 -

wherein: R¹ is C₈-C₂₄ alkyl or alkenyl;

R² is -H, C₁-C₄ alkyl, phenyl, or -CH₂COOM;

R³ is -C -R⁴₂; n is from 1 to 4; R⁴ is H, C₁-C₆ alkyl, or C₁-C₆ alkylester; and M is H or a cation, or a mixture thereof.

5

4. A shampoo composition as in Claim 3, wherein said N-acyl amino acid surfactant is an N-acyl sarcosinate surfactant.

5. A shampoo composition as in Claim 1, 2, 3, or 4, wherein said composition comprises from 5% to 15% of said alkyl ethoxylated sulfate, from 1.5% to 10% of said amphoteric surfactant, and from 0.5% to 5% of said N-acyl amino acid surfactant.

10

6. A shampoo composition as in Claim 1, 2, 3, 4, or 5 wherein a 30% aqueous solution of said amphoteric surfactant has an absorbance value at 440nm in a 1.0 path length of no greater than 0.035.

15

7. A shampoo composition as in Claim 1, 2, 3, 4, 5, or 6 wherein said absorbance value of said shampoo is no greater than 0.025 after 30 days at 38°C.

20

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 A61K7/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	US, A, 5 227 156 (R. S. WIESE) 13 July 1993 see the whole document ---	1-4
X	EP, A, 0 229 690 (THE PROCTER & GAMBLE COMPANY) 22 July 1987 see the whole document ---	1-5
X	EP, A, 0 150 323 (KAO CORPORATION) 7 August 1985 see the whole document ---	1, 5
A	EP, A, 0 356 784 (MITSUBISHI PETROCHEMICAL) 7 March 1990 -----	1-7

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "A" document member of the same patent family

Date of the actual completion of the international search

28 January 1994

Date of mailing of the international search report

09.02.94

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Sierra Gonzalez, M